


Boronic Acid for the Traceless Delivery of Proteins into Cells

Kristen A. Andersen,^{†,⊥} Thomas P. Smith,^{‡,⊥} Jo E. Lomax,[§] and Ronald T. Raines^{*,‡,||}[†]Graduate Program in Molecular and Cellular Pharmacology, University of Wisconsin—Madison, 1300 University Avenue, Madison, Wisconsin 53706, United States[‡]Department of Chemistry, University of Wisconsin—Madison, 1101 University Avenue, Madison, Wisconsin 53706, United States[§]Graduate Program in Cellular and Molecular Biology, University of Wisconsin—Madison, 1525 Linden Drive, Madison, Wisconsin 53706, United States^{||}Department of Biochemistry, University of Wisconsin—Madison, 433 Babcock Drive, Madison, Wisconsin 53706, United States

S Supporting Information

ABSTRACT: The use of exogenous proteins as intracellular probes and therapeutic agents is in its infancy. A major hurdle has been the delivery of native proteins to an intracellular site of action. Herein, we report on a compact delivery vehicle that employs the intrinsic affinity of boronic acids for the carbohydrates that coat the surface of mammalian cells. In the vehicle, benzoxaborole is linked to protein amino groups via a “trimethyl lock.” Immolation of this linker is triggered by cellular esterases, releasing native protein. Efficacy is demonstrated by enhanced delivery of green fluorescent protein and a cytotoxic ribonuclease into mammalian cells. This versatile strategy provides new opportunities in chemical biology and pharmacology.



The delivery of proteins and other macromolecules to an intracellular site is made difficult by cellular membranes.¹ Extensive efforts have led to the development of effective delivery systems that invoke cell-penetrating peptides,^{2–6} antibodies,⁷ ligands for natural receptors,⁸ dendrimers,⁹ functionalized polymers,^{10,11} liposomes,¹² or nanoparticles^{13,14} and even enable targeting to subcellular organelles.^{15,16} Nonetheless, extant strategies often inflict proteins with low biological activity,^{11,17–19} instability in a physiological context,^{20,21} or immunogenicity *in vivo*,^{22,23} or employ vehicles that are recalcitrant to biodegradation.²⁴

Boronic acids are physiologically benign Lewis acids that react spontaneously and reversibly with 1,2- and 1,3-diols to form five- and six-membered cyclic boronic esters, respectively.^{25,26} The dynamic covalent bonding of boronic acids/esters can facilitate the delivery of cargo into cells, which are coated with a diol-rich glycocalyx. To exploit that attribute, polymers, nanoparticles, and noncovalent assemblies have been decorated with phenylboronic acid and other arylboronic acids.^{27,28}

Recently, we showed that boronic acids can be advantageous when conjugated *directly* to a protein.¹⁸ The ensuing formation of transient boronate esters with the glycocalyx enhances cellular delivery. To date, this approach has relied on the irreversible modification of a target protein, with ensuing compromises. For example, our previous work installed boronic acids by the irreversible amidation of enzymic carboxyl groups, which led to the loss of 83% of catalytic activity.¹⁸ An ideal delivery system based on boronic acids (or, indeed, any moiety) conveys native cargo, and is thus “traceless.”

We sought to use a boronic acid and an immolative linker to promote the delivery of native proteins into a cell. As a boronic acid, we chose 2-hydroxymethylphenylboronic acid (benzoxaborole), which has higher affinity than does phenylboronic acid for the glycopyranosides that are abundant in the glycocalyx.^{18,28,29} As an immolative linker, we chose the *o*-hydroxydihydrocinnamic acid derivative known as the trimethyl lock (TML). After being triggered, the TML exhibits extremely high lactonization rates to release a cargo of interest (Scheme 1).^{30–34} The TML has been used for a wide variety of applications in chemistry and pharmacology³⁵ but not as an immolative linker on a protein. We chose ester hydrolysis as the means to trigger lactonization of the TML, as esterases are abundant inside, but not outside, of human cells^{36–38} and underlie the action of numerous prodrugs.³⁹ We equipped our TML scaffold with an *N*-hydroxysuccinimide ester for chemoselective conjugation to amino groups,¹⁷ such as those at the N terminus and on the side chain of lysine residues, which have a ~6% abundance in proteins.⁴⁰ Thus, our delivery vehicle (B-TML–NHS ester) has three modules: benzoxaborole, an esterase-activated TML linker, and an NHS ester (Figure 1A).

We synthesized B-TML–NHS ester convergently in 10 steps by extending a known procedure.⁴¹ Then, we characterized its ability to enhance the cellular internalization of green fluorescent protein (GFP) (Scheme 2), which has distinctive fluorescence and an inability to enter mammalian cells.⁴²

Received: August 1, 2015

Accepted: December 2, 2015

Published: December 2, 2015

Scheme 1




Figure 1. Cellular internalization of B-TML-labeled GFP. (A) Structures of B-TML-NHS ester and Ac-TML-NHS ester. Ellipses denote the three distinct modules within B-TML-NHS ester. (B) MALDI-TOF mass spectra of B-TML-GFP (green), conjugated to ~ 3 boronic acid moieties per molecule and the same protein after exposure to CHO K1 cell lysate and purification (gray). Expected m/z : GFP, 29361; each B-TML moiety, 450. (C) Flow cytometry analysis of CHO K1 cells incubated with 10 μ M unlabeled GFP, GFP labeled with a control vehicle (Ac-TML), or GFP labeled with the boronate vehicle (B-TML) for 4 h ($p < 0.0001$). (D) Confocal microscopy of CHO K1 cells grown as in panel C. Cells were stained with WGA-594 (red) and Hoechst 33342 (blue). Scale bars: 10 μ m.

Overnight incubation at ambient temperature with 100-fold excess of B-TML-NHS ester in 3:1 PBS/acetonitrile yielded 3 ± 1 labels per protein (Figures 1B and S1). The number of labels in the B-TML-GFP conjugate did not decrease after a

month of storage in PBS (Figure S2), consistent with the stability observed for other TML conjugates.^{43–45} Labeling was, however, “bioreversible.” Incubation with a lysate from Chinese hamster ovary (CHO) K1 cells removed all of the labels from B-TML-GFP (Figure 1B).

Next, we compared the uptake of B-TML-GFP and unlabeled GFP by CHO K1 cells. After a 4-h incubation, we observed a dramatic increase in the cellular uptake of B-TML-GFP (Figure 1C). The fluorescence in microscopy images was largely punctate, suggesting that B-TML-GFP was taken up *via* an endosomal pathway (Figure 1D). Co-localization of this bright punctate staining with a stain for transferrin was consistent with this conclusion (Figure S3). After a 24-h incubation, the fluorescence intensity had increased significantly beyond the level measured at 4 h, indicating that the conjugate was stable in the presence of serum, which was a component of the cell-culture medium (Figure S4).

To confirm that the boronic acid moiety was responsible for the difference in cellular entry, we performed two control experiments. First, we modified GFP with a vehicle (Ac-TML-NHS ester) that lacks the benzoxaborole functionality (Figure 1A), yielding a level of labeling similar to that from B-TML-NHS ester (Figure S1). When incubated with cells for 4 h, Ac-TML-GFP was taken up comparably to unlabeled GFP rather than to B-TML-GFP (Figure 1C and D). These data indicate that the enhanced delivery upon treatment with B-TML-NHS ester is not due to the mere modification of lysine residues or to interactions with the TML portion of B-TML. Next, we repeated the cellular uptake experiments with B-TML-GFP in the presence of fructose, which has a K_a of 336 M^{-1} for benzoxaborole.¹⁸ We observed a significant decrease in GFP uptake in the presence of fructose, apparent with both confocal microscopy and flow cytometry (Figure 2A and B). Again, these data indicate the boronic acid portion of B-TML-GFP as being responsible for cellular uptake.

Finally, we sought to test the efficacy of B-TML as a delivery vehicle to the cytosol. To do so, we employed an enzymic cytotoxin—the G88R variant of ribonuclease A, which must reach cytosolic RNA to manifest its toxic activity.^{46,47} After labeling the ribonuclease by the same procedure used to label GFP, we observed an average of 1.6 ± 0.7 labels per molecule of protein (Figure S5). This lower labeling is consistent with GFP (19 lysine residues) having more amino groups than does the ribonuclease (12 lysine residues). Again, we found that the labeling was bioreversible, as incubation with a CHO K1 cell lysate removed all of the labels (Figure S6). Finally, we assayed the ability of B-TML-ribonuclease and unlabeled ribonuclease to inhibit the proliferation of K-562 cells, which are derived from a human myelogenous leukemia line. We found that the pendant boronic acids resulted in a decrease in the IC_{50} value (Figure 3), indicative of more cytotoxin reaching the cytosol.

We conclude that covalent modification of proteins with B-TML-NHS ester can increase their ability to enter mammalian cells. Importantly, this modification is bioreversible. The

Scheme 2





Figure 2. Effect of fructose on the cellular internalization of B-TML-labeled GFP. (A) Confocal microscopy of B-TML-GFP (10 μ M) preincubated with PBS or 175 mM fructose for 30 min, then used to treat CHO K1 cells for 4 h. Cells were stained with WGA-594 (red) and Hoechst 33342 (blue). Scale bars: 20 μ m. (B) Flow cytometry analysis of CHO K1 cells treated as in panel A ($p < 0.01$).




Figure 3. Effect of B-TML-labeling on the inhibition of K-562 cell proliferation by a ribonuclease. Unlabeled G88R ribonuclease A, $IC_{50} = 6.4 \pm 0.1 \mu$ M; B-TML-labeled G88R ribonuclease A, $IC_{50} = 3.5 \pm 0.8 \mu$ M. Each data point represents the mean \pm SE for three separate experiments, each performed in duplicate.

irreversible modification of a protein can compromise its utility.^{11,17–23} In contrast, endogenous cellular enzymes revert modification with B-TML.

The bioreversibility of our delivery vehicle provides new opportunities. The sulfhydryl groups of cysteine residues have

long been used for this purpose because their mixed disulfides suffer reduction within the cytosol.^{48,49} Recently, we found that appropriately tuned diazo compounds can esterify protein carboxyl groups, providing a second type of bioreversible modification.^{50,51} In this work, we report on a bioreversible modification of protein amino groups that is distinct from others^{52–54} in its being removable by an enzyme-catalyzed reaction. With its traceless utility in promoting cellular uptake, B-TML-NHS ester provides new opportunities in chemical biology and pharmacology.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acschembio.5b00966.

Procedures for the synthesis of B-TML-NHS ester and Ac-TML-NHS ester, their use in protein modification, analyses of cellular internalization, and Figures S1–S6 (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: rtraines@wisc.edu.

Author Contributions

[†]These authors contributed equally to this work and are listed alphabetically.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to I. Ç. Tanrikulu (University of Wisconsin—Madison) for technical advice. K.A.A. was supported by a predoctoral fellowship from the PhRMA Foundation and Molecular and Cellular Pharmacology Training Grant T32 GM008688 (National Institutes of Health). J.E.L. was supported by a National Science Foundation Graduate Research Fellowship. This work was supported by grants R01 GM044786 and R01 CA073808 (National Institutes of Health) and made use of the National Magnetic Resonance Facility at Madison, which is supported by Grant P41 GM103399 (National Institutes of Health), and the University of Wisconsin Carbone Cancer Center, which is supported by Grant P30 CA014520 (National Institutes of Health).

REFERENCES

- (1) Fu, A., Tang, R., Hardie, J., Farkas, M. E., and Rotello, V. M. (2014) Promises and pitfalls of intracellular delivery of proteins. *Bioconjugate Chem.* 25, 1602–1608.
- (2) Fonesca, S. B., Pereira, M. P., and Kelley, S. O. (2009) Recent advances in the use of cell-penetrating peptides for medical and biological applications. *Adv. Drug Delivery Rev.* 61, 953–964.
- (3) Milletti, F. (2012) Cell-penetrating peptides: Classes, origin, and current landscape. *Drug Discovery Today* 17, 850–860.
- (4) Bechara, C., and Sagan, S. (2013) Cell-penetrating peptides: 20 Years later, where do we stand? *FEBS Lett.* 587, 1693–1702.
- (5) Brock, R. (2014) The uptake of arginine-rich cell-penetrating peptides: Putting the puzzle together. *Bioconjugate Chem.* 25, 863–868.
- (6) Di Pisa, M., Chassaing, G., and Swiecicki, J.-M. (2015) When cationic cell-penetrating peptides meet hydrocarbons to enhance in-cell cargo delivery. *J. Pept. Sci.* 21, 356–369.
- (7) Sievers, E. L., and Senter, P. D. (2013) Antibody–drug conjugates in cancer therapy. *Annu. Rev. Med.* 64, 15–29.
- (8) Srinivasarao, M., Galliford, C. V., and Low, P. S. (2015) Principles in the design of ligand-targeted cancer therapeutics and imaging agents. *Nat. Rev. Drug Discovery* 14, 203–219.
- (9) Cloninger, M. J. (2002) Biological applications of dendrimers. *Curr. Opin. Chem. Biol.* 6, 742–748.
- (10) Chu, D. S. H., Schellinger, J. G., Shi, J., Convertine, A. J., Stayton, P. S., and Pun, S. H. (2012) Application of living free radical polymerization for nucleic acid delivery. *Acc. Chem. Res.* 45, 1089–1099.
- (11) Gong, Y., Leroux, J. C., and Gauthier, M. A. (2015) Releasable conjugation of polymers to proteins. *Bioconjugate Chem.* 26, 1172–1181.
- (12) Allen, T. M., and Cullis, P. R. (2013) Liposomal drug delivery systems: From concept to clinical applications. *Adv. Drug Delivery Rev.* 65, 36–48.
- (13) Chou, L. Y. T., Ming, K., and Chan, W. C. W. (2011) Strategies for the intracellular delivery of nanoparticles. *Chem. Soc. Rev.* 40, 233–245.
- (14) Kumari, P., Ghosh, B., and Biswas, S. (2015) Nanocarriers for cancer-targeted drug delivery. *J. Drug Target.* 10, 1–13.
- (15) Rin Jean, S., Tulumello, D. V., Wisnovsky, S. P., Lei, E. K., Pereira, M. P., and Kelley, S. O. (2014) Molecular vehicles for mitochondrial chemical biology and drug delivery. *ACS Chem. Biol.* 9, 323–333.
- (16) Field, L. D., Delehanty, J. B., Chen, Y. C., and Medintz, I. L. (2015) Peptides for specifically targeting nanoparticles to cellular organelles: Quo vadis? *Acc. Chem. Res.* 48, 1380–1390.
- (17) Baslé, E., Joubert, N., and Pucheault, M. (2010) Protein chemical modification on endogenous amino acids. *Chem. Biol.* 17, 213–227.
- (18) Ellis, G. A., Palte, M. J., and Raines, R. T. (2012) Boronate-mediated biologic delivery. *J. Am. Chem. Soc.* 134, 3631–3634.
- (19) Boutureira, O., and Bernardes, G. J. (2015) Advances in chemical protein modification. *Chem. Rev.* 115, 2174–2195.
- (20) Fuchs, S. M., and Raines, R. T. (2005) Polyarginine as a multifunctional fusion tag. *Protein Sci.* 14, 1538–1544.
- (21) Fuchs, S. M., Rutkoski, T. J., Kung, V. M., Groeschl, R. T., and Raines, R. T. (2007) Increasing the potency of a cytotoxin with an arginine graft. *Protein Eng., Des. Sel.* 20, 505–509.
- (22) De Groot, A. S., and Scott, D. W. (2007) Immunogenicity of protein therapeutics. *Trends Immunol.* 28, 482–490.
- (23) Pisal, D. S., Kosloski, M. P., and Balu-Iyer, S. V. (2010) Delivery of therapeutic proteins. *J. Pharm. Sci.* 99, 2557–2575.
- (24) Onoue, S., Yamada, S., and Chan, H. K. (2014) Nanodrugs: Pharmacokinetics and safety. *Int. J. Nanomed.* 9, 1025–1037.
- (25) Bosch, L. L., Fyles, T. M., and James, T. D. (2004) Binary and ternary phenylboronic acid complexes with saccharides and Lewis bases. *Tetrahedron* 60, 11175–11190.
- (26) Peters, J. A. (2014) Interactions between boric acid derivatives and saccharides in aqueous media: Structures and stabilities of resulting esters. *Coord. Chem. Rev.* 268, 1–22.
- (27) Ma, R., and Shi, L. (2014) Phenylboronic acid-based glucose-responsive polymeric nanoparticles: Synthesis and applications in drug delivery. *Polym. Chem. S.* 1503–1518.
- (28) Adamczyk-Woźniak, A., Borys, K. M., and Sporyński, A. (2015) Recent developments in the chemistry and biological applications of benzoxaboroles. *Chem. Rev.* 115, 5224–5247.
- (29) Bérubé, M., Dowlut, M., and Hall, D. G. (2008) Benzoboroxoles as efficient glycopyranoside-binding agents in physiological conditions: Structure and selectivity of complex formation. *J. Org. Chem.* 73, 6471–6479.
- (30) Milstien, S., and Cohen, L. A. (1970) Concurrent general-acid and general-base catalysis of esterification. *J. Am. Chem. Soc.* 92, 4377–4382.
- (31) Danforth, C., Nicholson, A. W., James, J. C., and Loudon, G. M. (1976) Steric acceleration of lactonization reactions: An analysis of stereopopulation control. *J. Am. Chem. Soc.* 98, 4275–4281.
- (32) Wang, B., Nicolaou, M. G., Liu, S., and Borchardt, R. T. (1996) Structural analysis of a facile lactonization system facilitated by a trimethyl lock. *Bioorg. Chem.* 24, 39–49.
- (33) Nicolaou, M. G., Wolfe, J. L., Schowen, R. L., and Borchardt, R. T. (1996) Facilitated intramolecular conjugate addition of amides of 3-(3',6'-dioxo-2",4"-dimethyl-1",4"-cyclohexadienyl)-3,3-dimethylpropionic acid. 2. Kinetics of degradation. *J. Org. Chem.* 61, 6633–6638.
- (34) Jung, M. E., and Piizzi, G. (2005) *gem*-Disubstituent effect: Theoretical basis and synthetic applications. *Chem. Rev.* 105, 1735–1766.
- (35) Levine, M. N., and Raines, R. T. (2012) Trimethyl lock: A trigger for molecular release in chemistry, biology, and pharmacology. *Chem. Sci.* 3, 2412–2420.
- (36) Liederer, B. M., and Borchardt, R. T. (2006) Enzymes involved in the bioconversion of ester-based prodrugs. *J. Pharm. Sci.* 95, 1177–1195.
- (37) Lavis, L. D. (2008) Ester bonds in prodrugs. *ACS Chem. Biol.* 3, 203–206.
- (38) Fukami, T., and Yokoi, T. (2012) The emerging role of human esterases. *Drug Metab. Pharmacokinet.* 27, 466–477.
- (39) Testa, B., and Mayer, J. M. (2003) *Hydrolysis in Drug and Prodrug Metabolism: Chemistry, Biochemistry, and Enzymology*, Verlag Helvetica Chimica Acta, Zürich, Switzerland.
- (40) McCaldon, P., and Argos, P. (1988) Oligopeptide biases in protein sequences and their use in predicting protein coding regions in nucleotide sequences. *Proteins: Struct., Funct., Genet.* 4, 99–122.
- (41) Greenwald, R. B., Choe, Y. H., Conover, C. D., Shum, K., Wu, D., and Royzen, M. (2000) Drug delivery systems based on trimethyl lock lactonization: Poly(ethylene glycol) prodrugs of amino-containing compounds. *J. Med. Chem.* 43, 475–487.
- (42) Fuchs, S. M., and Raines, R. T. (2007) Arginine grafting to endow cell permeability. *ACS Chem. Biol.* 2, 167–170.
- (43) Lavis, L. D., Chao, T.-Y., and Raines, R. T. (2006) Fluorogenic label for biomolecular imaging. *ACS Chem. Biol.* 1, 252–260.
- (44) Johnson, R. J., Chao, T.-Y., Lavis, L. D., and Raines, R. T. (2007) Cytotoxic ribonucleases: The dichotomy of Coulombic forces. *Biochemistry* 46, 10308–10316.
- (45) Turcotte, R. F., Lavis, L. D., and Raines, R. T. (2009) Onconase cytotoxicity relies on the distribution of its positive charge. *FEBS J.* 276, 3846–3857.
- (46) Leland, P. A., Schultz, L. W., Kim, B.-M., and Raines, R. T. (1998) Ribonuclease A variants with potent cytotoxic activity. *Proc. Natl. Acad. Sci. U. S. A.* 95, 10407–10412.
- (47) Lomax, J. E., Eller, C. H., and Raines, R. T. (2012) Rational design and evaluation of mammalian ribonuclease cytotoxins. *Methods Enzymol.* 502, 273–290.
- (48) Gilbert, H. F. (2006) Molecular and cellular aspects of thiol–disulfide exchange. *Adv. Enzymol.* 63, 69–172.
- (49) Robbins, P. B., Oliver, S. F., Sheu, S. M., Goodnough, J. B., Wender, P., and Khavari, P. A. (2002) Peptide delivery to tissues via

reversibly linked protein transduction sequence. *BioTechniques* 33, 190–194.

(50) McGrath, N. A., Andersen, K. A., Davis, A. K. F., Lomax, J. E., and Raines, R. T. (2015) Diazo compounds for the bioreversible esterification of proteins. *Chem. Sci.* 6, 752–755.

(51) Mix, K. A., and Raines, R. T. (2015) Optimized diazo scaffold for protein esterification. *Org. Lett.* 17, 2358–2361.

(52) Dixon, H. B. F., and Perham, R. N. (1968) Reversible blocking of amino groups with citraconic anhydride. *Biochem. J.* 109, 312–314.

(53) Wang, M., Sun, S., Neufeld, C. I., Perez-Ramirez, B., and Xu, Q. (2014) Reactive oxygen species-responsive protein modification and its intracellular delivery for targeted cancer therapy. *Angew. Chem., Int. Ed.* 53, 13444–13448.

(54) Cal, P. M., Frade, R. F., Cordeiro, C., and Gois, P. M. (2015) Reversible lysine modification on proteins by using functionalized boronic acids. *Chem. - Eur. J.* 21, 8182–8187.